

Table 2a. AIDS diagnoses, by year of diagnosis and selected characteristics, 2006–2009 and cumulative—United States

	2006			2007			2008			2009			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Age at diagnosis (yr)														
<13	37	39	0.1	29	31	0.1	35	40	0.1	10	13	0.0	9,369	9,448
13–14	73	78	0.9	74	80	1.0	51	58	0.7	39	58	0.7	1,262	1,321
15–19	376	398	1.9	416	449	2.1	429	480	2.2	376	484	2.2	6,910	7,214
20–24	1,548	1,637	7.7	1,750	1,880	8.9	1,675	1,869	8.8	1,667	2,095	9.7	41,628	42,920
25–29	3,176	3,353	16.3	3,135	3,378	16.1	3,064	3,428	16.0	2,753	3,476	16.0	126,954	129,639
30–34	4,048	4,279	22.0	3,805	4,096	21.2	3,716	4,147	21.2	3,227	4,043	20.3	210,270	214,149
35–39	5,911	6,254	29.8	5,317	5,741	27.3	4,776	5,367	25.7	3,888	4,893	23.8	229,584	234,575
40–44	6,760	7,159	32.1	6,181	6,680	30.6	5,540	6,222	29.1	4,502	5,689	27.1	188,102	193,237
45–49	5,236	5,543	24.4	5,307	5,731	25.2	4,885	5,518	24.2	4,303	5,466	23.9	122,228	126,380
50–54	3,405	3,608	17.7	3,340	3,599	17.2	3,274	3,660	17.1	3,138	3,983	18.3	69,691	72,327
55–59	1,901	2,013	11.1	1,855	1,998	11.0	1,877	2,091	11.3	1,723	2,191	11.5	37,604	39,025
60–64	900	954	7.2	909	979	6.8	965	1,075	7.1	805	1,010	6.4	19,947	20,633
≥65	788	836	2.2	732	791	2.1	721	800	2.1	669	846	2.1	17,165	17,743
Race/ethnicity														
American Indian/Alaska Native	138	145	6.4	132	141	6.1	161	177	7.6	127	155	6.6	3,599	3,700
Asian ^c	395	422	3.3	416	454	3.5	434	489	3.7	329	429	3.1	7,995	8,324
Black/African American	16,357	17,300	47.4	15,926	17,171	46.5	15,226	17,065	45.7	13,196	16,741	44.4	452,985	466,351
Hispanic/Latino ^d	6,652	7,065	16.1	6,390	6,912	15.2	5,962	6,701	14.3	5,285	6,719	13.9	184,719	190,263
Native Hawaiian/Other Pacific Islander	47	49	11.7	48	51	12.0	37	41	9.2	40	50	11.2	810	839
White	9,924	10,484	5.3	9,331	10,050	5.0	8,640	9,668	4.8	7,595	9,467	4.7	418,074	426,102
Multiple races	646	686	16.5	607	656	15.3	548	616	13.9	528	686	15.1	12,229	12,726
Transmission category														
Male adult or adolescent														
Male-to-male sexual contact	13,006	16,517	—	12,621	16,521	—	11,871	16,469	—	10,773	17,005	—	473,724	529,908
Injection drug use	2,743	3,853	—	2,361	3,505	—	2,061	3,303	—	1,523	3,012	—	160,614	186,318
Male-to-male sexual contact and injection drug use	1,595	1,957	—	1,394	1,798	—	1,264	1,706	—	1,000	1,580	—	70,586	77,213
Heterosexual contact ^e	2,829	3,952	—	2,607	3,882	—	2,552	3,949	—	2,148	3,832	—	56,074	72,183
Other ^f	4,827	194	—	4,964	165	—	5,072	185	—	4,741	158	—	96,383	12,744
Subtotal	25,000	26,473	22.0	23,947	25,871	21.3	22,820	25,612	20.9	20,185	25,587	20.6	857,381	878,366

Table 2a. AIDS diagnoses, by year of diagnosis and selected characteristics, 2006–2009 and cumulative—United States (cont)

	2006			2007			2008			2009			Cumulative ^b	
	No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Estimated ^a		No.	Est. No. ^a
		No.	Rate		No.	Rate		No.	Rate		No.	Rate		
Female adult or adolescent														
Injection drug use	1,599	2,469	—	1,491	2,392	—	1,221	2,141	—	900	1,930	—	70,856	87,126
Heterosexual contact ^e	4,580	6,980	—	4,355	6,955	—	3,965	6,824	—	3,241	6,561	—	96,111	126,637
Other ^f	2,943	190	—	3,027	184	—	2,967	137	—	2,764	155	—	46,995	7,032
Subtotal	9,122	9,639	7.7	8,873	9,531	7.5	8,153	9,102	7.1	6,905	8,647	6.7	213,962	220,795
Child (<13 yrs at diagnosis)														
Perinatal	34	36	—	28	30	—	31	35	—	9	12	—	8,568	8,640
Other ^g	3	3	—	1	1	—	4	5	—	1	1	—	801	807
Subtotal	37	39	0.1	29	31	0.1	35	40	0.1	10	13	0.0	9,369	9,448
Region of residence														
Northeast	8,742	9,369	17.1	8,298	9,082	16.5	6,986	8,064	14.6	5,807	8,171	14.8	330,979	340,357
Midwest	3,903	4,154	6.3	3,675	4,006	6.0	3,687	4,218	6.3	3,435	4,394	6.6	112,560	116,029
South	15,718	16,453	15.1	15,371	16,383	14.8	15,003	16,506	14.7	13,188	15,806	13.9	419,910	430,141
West	5,796	6,174	9.0	5,506	5,964	8.5	5,332	5,967	8.4	4,670	5,875	8.2	217,265	222,083
Total^h	34,159	36,151	12.1	32,850	35,434	11.7	31,008	34,755	11.4	27,100	34,247	11.2	1,080,714ⁱ	1,108,611

^a Estimated numbers resulted from statistical adjustment that accounted for reporting delays and missing risk-factor information, but not for incomplete reporting. Rates are per 100,000 population. Rates are not calculated by transmission category because of the lack of denominator data.

^b From the beginning of the epidemic through 2009.

^c Includes Asian/Pacific Islander legacy cases (see Technical Notes).

^d Hispanics/Latinos can be of any race.

^e Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^f Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

^g Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^h Because column totals for estimated numbers were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

ⁱ Includes persons of unknown race/ethnicity.